

*Editor's note: This English translation was done by a third party. The original document can be accessed [here](#).*

**Federal Agency for Medicines and Health Products**  
**Finasteride EG Leaflet**  
**Leaflet: Information for the Consumer**  
**Finasteride EG 1 mg film-coated tablets**

**Read the entire leaflet carefully before you start taking the medicine because it contains important information for you.**

- Keep this leaflet. You might need to read it again
- If you have any additional questions ask your doctor or pharmacist
- This medicine is prescribed only to you. Do not recommend it to other people. It might harm them regardless of the similarity of their symptoms to yours.
- If you experience any adverse reactions, inform your doctor or pharmacist. This includes all adverse reactions not described in this leaflet. Check point 4.

**What this leaflet contains:**

1. What is Finasteride EG 1 mg film-coated tablets and what is it used for
2. What do you have to know before you take Finasteride EG 1 mg film-coated tablets
3. How to take Finasteride EG 1 mg film-coated tablets
4. Possible adverse reactions
5. How to store Finasteride EG 1 mg film-coated tablets
6. Package content and additional information

**1. What is Finasteride EG 1 mg film-coated tablets and what is it used for**

Finasteride EG 1 mg film-coated tablets contains finasteride. Finasteride belongs to a group of medicines called 5 $\alpha$ -reductase inhibitors.

Your doctor prescribed you Finasteride EG 1 mg film-coated tablets because you have male pattern baldness (also known as androgenic alopecia). Finasteride EG 1 mg film-coated protects against further hair loss in men. Men with mild to medium, but not full baldness, can benefit by the usage of Finasteride EG 1 mg film-coated tablets. Finasteride EG 1 mg film-coated tablets block an important enzyme (type II 5 $\alpha$ -reductase) that participates in the hair follicle regulation.

Finasteride EG 1 mg film-coated tablets specifically decrease the dihydrotestosterone (DHT) level in the scalp, the main reason for male pattern baldness. In this way Finasteride EG 1 mg film-coated tablets help to stop the baldness process and to prevent further hair loss.

**2. What you must know before you take Finasteride EG 1 mg film-coated tablets**

Do not take Finasteride EG 1 mg:

- If you are allergic to finasteride or any of the other ingredients in this medicine (they are listed in point 6).
- If you are a child or youth.

If you are a woman (see section below: "Pregnancy, breastfeeding and fertility").

- During the clinical trials, it was concluded that Finasteride EG 1 mg film-coated tablets is not effective in hair loss treatment (androgenic alopecia) in women.

- If you are already taking finasteride or any other 5 $\alpha$ -reductase inhibitor to treat benign prostatic hyperplasia or any other condition.

### **Warnings and precautions**

- Talk to your doctor or pharmacist before you take Finasteride EG 1 mg film-coated tablets.
- Before starting treatment and during the patients' therapy with finasteride, determination for the prostate-specific antigen (PSA) in the serum must be performed.
- Finasteride may affect male fertility. Patients that are planning to have a child must consider stopping the treatment.

### **Mood swings and depression**

Mood swings such as depressive mood, depression and, rarely, suicide thoughts, have been reported in patients treated with Finasteride EG 1 mg film-coated tablets. If you have any of these symptoms, stop taking Finasteride EG 1 mg film-coated tablets and contact your doctor as soon as possible for additional medical help.

### **Other medicine and Finasteride EG 1 mg film-coated tablets**

Inform your doctor or pharmacist if you take or recently have been taking or if it is possible that you are taking other medicine. Finasteride EG 1 mg film-coated tablets usually does not interact with other medicines.

There is no available information regarding the simultaneous usage of Finasteride EG 1 mg film-coated tablets and topical minoxidil (applied on the skin) in male pattern baldness. This combination is not recommended.

### **Finasteride EG 1 mg film-coated tablets intake with food and beverages.**

Finasteride EG 1 mg can be taken with or without food.

### **Pregnancy, breastfeeding and fertility**

Finasteride EG 1 mg film-coated tablets are intended only for men. Women, who are pregnant or have the possibility of becoming pregnant, must not touch broken tablets that contains finasteride. If finasteride is absorbed by the skin or is being taken orally by a pregnant woman with a male baby, the baby might be born with deformed genitals.

When the intimate partner of the patient is pregnant or has the possibility of becoming pregnant, it is recommended that the patient reduce her contact to his semen to a minimum (for example, with the use of a condom) or to stop the treatment with finasteride.

### **Driving and working with machines**

There is no data suggesting that Finasteride EG 1 mg film-coated tablets influences the ability to drive or work with machines.

### **Important information regarding some ingredients of Finasteride EG 1 mg film-coated tablets**

This medical product contains lactose. If your doctor has told you that you have intolerance to some sugars, contact them before beginning to take this medicine.

### **3. How to take Finasteride EG 1 mg film-coated tablets**

Always take the medication exactly the way your doctor has told you. If you are unsure about something, ask your doctor or pharmacist.

The recommended dose is one tablet of Finasteride EG 1 mg daily (that equals 1 mg finasteride). The film-coated tablets can be taken on empty stomach, or with food. The film-coated tablets must be swallowed whole and they must not be divided or broken.

- Finasteride EG 1 mg film-coated tablets will not have faster or stronger effect if they are taken more than once daily.
- Male pattern baldness is a condition that develops over a long period of time. Most often it is necessary to use daily for 3 to 6 months before noticing any hair growth increase or hair loss decrease.
- Your doctor will help you to evaluate whether Finasteride EG 1 mg film-coated tablets have any effect. It is important to continue taking Finasteride EG 1 mg film-coated tablets as long as your doctor has prescribed to you.

#### **Patients with impaired liver functions**

There is no data for the usage of Finasteride EG 1 mg film-coated tablets in patients with impaired liver function.

#### **Patients with impaired kidney functions**

There is no need for dosage change in patients with impaired kidney functions.

#### **If you have taken more than the needed dose of Finasteride EG 1 mg film-coated tablets**

If you have taken more than the needed dose of Finasteride EG 1 mg film-coated tablets or anyone else has taken Finasteride EG 1 mg film-coated tablets, immediately contact the emergency ward of your nearest hospital. Show the doctor the rest of the tablets and your medicine package.

#### **If you have missed taking Finasteride EG 1 mg film-coated tablets**

If you forget to take a dose from Finasteride EG 1 mg film-coated tablets, just take the next dose as usual. Do not take double dose to compensate for the missed one.

#### **If you have stopped taking Finasteride EG 1 mg film-coated tablets**

In order to achieve maximum effect, continued use of Finasteride EG 1 mg film-coated tablets is recommended. If, meanwhile, you stop the treatment, in the next 9 to 12 months you will lose the hair that you have acquired.

If you have any questions regarding the usage of this medicine, ask your doctor or pharmacist.

#### **4. Possible adverse reactions**

As with every medicine, this medicine can cause adverse reactions despite the fact that not everyone experiences them. Some of the adverse reactions are temporary during the treatment or disappear when you stop the treatment.

**Stop Finasteride EG 1 mg film-coated tablets and immediately contact a doctor if you develop any of the following:**

- Allergic reaction symptoms, such as swelling of the lips, face, tongue, throat; experience difficulty in swallowing, rash (hives) and difficulty in breathing
- Depression (feeling of strong sadness and low self-esteem).

You must immediately inform your physician for any change in your breast tissue, such as lumps, breast enlargement, nipple secretion as this could be a sign of a serious disease, such as breast cancer.

#### **Uncommon (can affect between 1 and 100 people)**

- lower libido;
- erection difficulties;
- ejaculation problems, such as decrease in the ejected semen during sex
- depression

#### **Unknown frequency**

- allergic reaction, such as rash, itching, hives, lips and face swelling;
- breast pain and swelling;
- palpitations;
- testicle pain;
- increased heart rate;
- permanent erectile dysfunction after stopping the treatment;
- permanent decrease in libido after stopping the treatment;
- permanent ejaculation problems after stopping the treatment;
- there is data for infertility in men who have taken finasteride for a continued long period and have also had other risk factors that might have affected fertility. Normalization and improvement in the semen quality was reported after stopping the treatment with finasteride. Long-term clinical trials on finasteride's effect on men's fertility rate have not been done.
- Changes in the liver function that can be detected in blood tests
- Anxiety

#### **Reporting adverse reactions**

If you develop any side effects, talk to your doctor or pharmacist. This also applies to any adverse effects not mentioned in this leaflet. You can also report side effects directly.

In Belgium:

FAMHP  
Vigilance Division  
PO Box 97  
B-1000 Brussels Madou  
website: [www.afmps.be](http://www.afmps.be).

In Luxembourg:

Regional Pharmacovigilance Center of Nancy

E-mail: [crpv@chru-nancy.fr](mailto:crpv@chru-nancy.fr)

Tel.: (+33) 3 83 65 60 85 / 87

Fax: (+33) 3 83 65 61 33

Pharmacy and Medicines Division

Directorate of Health in Luxembourg

E-mail: [pharmacovigilance@ms.etat.lu](mailto:pharmacovigilance@ms.etat.lu)

Tel.: (+352) 247-85592

Fax: (+352) 247-95615

Link for the form: <http://www.sante.public.lu/fr/politiquesante/ministry-health/direction-health/div-pharmacy-medicines/index.html>.

By reporting side effects, you help provide more safety information for the drug.

## 5. How to store Finasteride EG 1 mg film-coated tablets

- keep out of reach of children
- this medicinal product does not require special storage conditions
- Do not use Finasteride EG 1 mg film-coated tablets if expired after the date marked on the packaging. The expiration day is the last day of the indicated month.
- These medicines must not be thrown down a drain or into regular trash bins. Ask your pharmacist how to dispose of the medicine that you are no longer using. This measure will help to keep the environment clean.

## 6. Package content and additional information

### What Finasteride EG 1 mg film-coated tablets contain

The active ingredient is finasteride. One film-coated tablet contains 1 mg finasteride.

Other ingredients are:

Tablet's core: lactose monohydrate, microcrystalline cellulose (E460); pregelatinised starch, sodium starch glycolate (type A); lauryl microglycerides, magnesium stearate (E570)

Film coat: Hypromellose (E464), titanium dioxide (E 171), macrogol 6000, red iron oxide (E172), yellow iron oxide (E172)

### What do Finasteride EG 1 mg film-coated tablets look like and what does the package contain:

Film-coated tablet.

Finasteride EG 1 mg are reddish-brown, round-shaped, with a sign "F1" from one side and without a sign on the other side.

Finasteride EG 1 mg tablets are packed in blisters aluminum-aluminum, in 28, 30, 84 or 98 package tablets.

## Marketing Authorization Holder

EG (Eurogenerics) SA  
Esplanade Heysel b22

1020 Brussels, Belgium

### **Manufacturers**

Kern Pharma, S.L.  
Venus 72, 08228 Terrassa  
Barcelona, Spain

Stada Arzneimittel AG  
Stadastraße 2-18 – 61118  
Bad Vilbel, Germany

Haupt Pharma Münster GmbH  
Schleebrüggekamp 15, 48159  
Münster, Germany

Stada Arzneimittel GesmbH  
Muthgasse 36/2, 1190  
Vienna, Austria

Eurogenerics SA  
Esplanade Heysel b22  
1020 Brussels, Belgium

PharmaCoDane Aps  
Marielundvej 46A  
2730 Herlev, Denmark

Siegfried Malta Ltd.  
HHF070 Hal Far Industrial Estate  
Hal Far BGG3000  
Malta

This medicinal product is authorized in the Member States of the European Economic Area under the following names:

Austria: Finasterid Stada 5 mg Filmtabletten

Belgium: Finasteride EG 5 mg film-coated tablets

Denmark: Finasterid "Stada"

Germany: Finasterid AL 5 mg Filmtabletten

Iceland: Finasterid Stada 5 mg filmuhúðuð tafla

Italy: Finasteride EG 5 mg rivestite coated tablets

Luxembourg: Finasteride EG 5 mg film-coated tablets

Poland: Finasterid Stada 5 mg tabletki powlekane

Sweden: Finasterid Stada 5 mg filmdragerad tablett

Marketing authorization numbers: BE294262 (plate) BE294271 (bottle)

Mode of delivery: via medical prescription.

The last date this leaflet was approved / revised is 03/2020 / 11/2019