

Editor's note: This English translation was done by a third party. The original report can be accessed [here](#).

**PACKAGE LEAFLET
INFORMATION FOR PATIENTS
Accord Finasteride 5 mg film-coated tablets**

Active ingredient: finasteride

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you. Keep the leaflet. You might want to read it again. If you have any further questions, ask your doctor or pharmacist.

This medicine has been prescribed for you personally. Do not pass it on to others. It can harm other people, even if they have the same symptoms as you. If you get any side effects, talk to your doctor or pharmacist. This also applies to side effects that are not listed in this leaflet. See section 4.

What is in this leaflet

1. What Accord Finasteride is and what it is used for
2. What you need to know before you take Accord Finasteride
3. How to take Accord Finasteride
4. What side effects are possible?
5. How to store Accord Finasteride
6. Contents of the pack and other information

1. WHAT ACCORD FINASTERIDE IS AND WHAT IT IS USED FOR

Accord Finasteride is used by men to treat and control

• Benign enlargement of the prostate (non-aggressive prostate growth).

Accord Finasteride belongs to a group of medicines called 5-alpha reductase inhibitors. It shrinks the prostate gland in men, improves urinary flow and other symptoms caused by BPH. Accord Finasteride reduces the risk of acute urinary retention and the need for surgery.

2. BEFORE YOU TAKE ACCORD FINASTERIDE

Accord Finasteride must not be taken if you

- are allergic to:
 - finasteride
 - or any of the other ingredients of this medicine (listed in section 6)
- are a woman
- are a youth or child under 18 years of age

If you are taking Accord Finasteride to treat an enlarged prostate, you should see a urologist.

Warnings and Precautions

Talk to your doctor before taking Accord Finasteride. Tell your doctor if any of the following apply to you:

- If you have difficulty emptying your bladder completely or have a strong void have decreased urinary flow Your doctor will examine you carefully before you start treatment with Accord Finasteride Rule out the possibility of another urinary tract disease.
- if you need a blood test for PSA, a determination of the prostate-specific antigen (PSA) Talk to your doctor or nurse about taking Accord Finasteride take as it may affect your test results.
- if you have impaired liver function

Women who are pregnant or may be pregnant should not use finasteride come into contact with Sandoz (for example via the semen or broken or crumbled tablets. See section Pregnancy and breast-feeding).

Mood changes and depression

In patients treated with Accord Finasteride, mood changes such as depressed mood, depression and, less commonly reported, suicidal thoughts. If you notice any of these symptoms, consult your doctor immediately.

Taking Accord Finasteride with other medicines

Tell your doctor or pharmacist if you are taking any other medicines are taking/using, have recently taken/used other medicines or plan to take/use other medicines. Accord Finasteride is not known to affect or be affected by any other medicines other medicines.

Pregnancy and breastfeeding period

Accord Finasteride should not be taken by women.

- If your sexual partner is pregnant or may be pregnant: The semen may contain traces of Accord Finasteride. That's why your sperm can go with you not come into contact with your partner, e.g. by using a condom.
- Women who are pregnant or may become pregnant should not use broken ones or crushed Accord Finasteride film-coated tablets.

If Accord Finasteride by a pregnant woman with a male fetus is ingested, the child may be born with malformed genitals will. The tablets are therefore film coated to prevent contact with finasteride to prevent Sandoz. There have been reports of infertility following the marketing of finasteride-containing products (infertility) and/or poor semen quality. Patients present other risk factors that may have contributed to infertility. A normalization or improvement in semen quality was observed after weaning finasteride reported. If you have any questions about this, please seeing your doctor.

Ability to drive and use machines There is no information that Accord Finasteride affects the ability to drive or use machines. Finsterid Sandoz contains lactose and sodium This medicine contains lactose. Please take Accord Finasterideerst afterwards Talk to your doctor if you know you have a suffer from sugar intolerance. 3 This medicinal product contains less than 1 mmol (23 mg) sodium per film-coated tablet, i.e. it is almost "sodium-free".

3. HOW TO TAKE ACCORD FINASTERIDE

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure. The recommended dose is:

- 1 tablet once a day type of administration

Swallow the film-coated tablet whole with a glass of water. You can with or without food can be taken, but always at the same time.

Duration of use

Your doctor will decide how long you should stop taking Accord Finasteride. Although improvement is often observed after a short period of time, it is necessary to continue treatment for at least 6 months. If you take more Accord Finasteride than you do should, contact your doctor or pharmacist immediately.

If you forget to take Accord Finasteride

If you forget to take a dose, do not double the dose. Continue taking the next dose as usual.

If you stop taking Accord Finasteride

Do not change or stop treatment without your doctor's permission. This can cause serious injury and impair the success of the treatment. If you have any further questions on the use of this medicine, please contact your doctor or pharmacist.

4. WHAT SIDE EFFECTS ARE POSSIBLE?

Like all medicines, this medicine can cause side effects, although these are not common have to appear to everyone. The most common side effects are the inability to get an erection and decreased sexual drive. These side effects occur in the early treatment phase appear and in most cases disappear in the further course of treatment. You should inform your doctor immediately if you notice any changes in your breast tissue such as lumps, pain or discharge from the nipples as this could be signs of a serious illness, such as breast cancer.

Allergic reaction

If you have an allergic reaction, stop taking it and let them know your doctor immediately. The signs of this can be:

- skin rash, itching or blistering under the skin (hives)
- swelling of your lips, tongue, throat and face

Side effects can occur with the following frequencies:

Common, may affect up to 1 in 10 people:

- inability to have an erection
- decreased sexual drive
- decreased sperm volume

Uncommon, may affect up to 1 in 100 people:

- tenderness to touch and/or breast enlargement
- ejaculation disorders
- depression

Very rare, may affect up to 1 in 10,000 people:

- secretion from the male breast
- breast lumps, which in some cases are surgically removed

Not known, frequency cannot be estimated from the available data:

- palpitations
- increased liver enzymes
- testicle pain
- inability to get an erection that persists after stopping treatment
- decreased sexual drive that persists after stopping treatment
- male infertility and/or poor sperm quality. It has been reported that sperm quality improves after stopping treatment returned to normal or improved.
- anxiety

Reporting side effects

If you get any side effects, talk to your doctor or pharmacist. This also applies to side effects not listed in this leaflet. You can also report side effects directly via the national reporting system:

Pharmaceutical Services
Ministry of Health
CY-1475 Nicosia
Fax: + 357 22608649
Website: www.moh.gov.cy/phs.

By reporting side effects, you can help provide more information be provided about the safety of this drug.

5. HOW TO STORE ACCORD FINASTERIDE

Keep this medicine out of the sight and reach of children. Do not use this medicine after the expiration date printed on the carton and blister after "Use by." The expiration date refers to the last day of the specified month. Do not store above 25°C. Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use, thus helping to protect the environment.

6. CONTENTS OF THE PACK AND OTHER INFORMATION

What Accord Finasteride contains

The active substance is finasteride. Each film-coated tablet contains 5 mg Finasteride. The other ingredients are: Docusate Sodium, Hypromellose, Indigo Carmine Salt (E132), Lactose Monohydrate, Magnesium Stearate, Microcrystalline Cellulose, Povidone K30, Propylene Glycol, Sodium Starch (Type A), Talc, Titanium Dioxide (E171).

What Finasteride Accord 1 mg tablets look like and contents of the pack

Film-coated tablet. Finasteride Accord 1 mg tablets are brown-red, round, biconvex, film-coated tablets marked "F1" on one side and blank on the other. Finasteride Accord 1 mg tablets are packed in aluminum-aluminum blisters in packs of 28, 30, 84 and 98 tablets.

Marketing Authorization Holder and Manufacturer: Accord Healthcare S.L.U. World Trade Center, Moll de Barcelona, s / n, Edifici Est 6^a planta, 08039 Barcelona, Spain

Manufacturer

Accord Healthcare Limited
Sage House, 319, Pinner Road
North Harrow, Middlesex, HA1 4HF
United Kingdom.

Accord Healthcare Polska Sp.z o.o.
Lutomierska 50,95-200 Pabianice
Poland

Accord Healthcare BV
Winthontlaan 200, 3526 KV
Utrecht, Netherlands

Laboratorios Normon SA
Ronda de Valdecarrizo 6,
28760 Tres Cantos
Madrid, Spain

This medicinal product is approved in the Member States of the European Economic Area (EEA) under the following names:

Estonia: Finasteride Accord 1 mg
Italy: Finasteride AHCL 1 mg
Malta: Finasteride 1 mg film-coated tablets
Spain: Finasterida Norman 1 mg compressed tablets with EFG film
United Kingdom: Finasteride 1 mg Film-coated Tablets
Bulgaria: Finasteride Accord 1 mg film-coated tablets
Cyprus: Finasteride Accord 1 mg film-coated tablets
Latvia: Finasteride Accord 1 mg apvalkotās tablets
Poland: Finahit, 1 mg, tabletki powlekane
Sweden: Finasteride Accord 1 mg filmdragerade tabletter
Netherlands: Finasteride Accord 1 mg filmomhulde tabletten
Portugal: FINOX 1 mg comprimidos revestidos per película

This brochure was last revised on 12/08/2020.